

# Veröffentlichungen Verbundprojekt „Organische Böden“

(Stand: 17.01.2018)

## Messung von Treibhausgasemissionen

Beetz, S., Liebersbach, H., Glatzel, S., Jurasinski, G., Buczko, U. & H. Höper, 2013: Effects of land use intensity on the full greenhouse gas balance in an atlantic peat bog, Biogeosciences 10: 1067–1082.

<http://www.biogeosciences.net/10/1067/2013/bg-10-1067-2013.html>

Beyer, C. & H. Höper, 2014: Greenhouse gas emissions from rewetted bog peat extraction sites and a Sphagnum cultivation site in Northwest Germany. Biogeosciences 12: 2101–2117.

<http://www.biogeosciences.net/12/2101/2015/bg-12-2101-2015.html>

Beyer, C., Liebersbach, H., & H. Höper, 2015: Multiyear greenhouse gas flux measurements on a temperate fen soil used for cropland or grassland. Journal of Plant Nutrition and Soil Science 178: 99–111, 2015.

<http://onlinelibrary.wiley.com/doi/10.1002/jpln.201300396/abstract>

Eickenscheidt T., Freibauer, A., Heinichen, J., Augustin, J. & M. Drösler, 2014: Short-term effects of biogas digestate and cattle slurry application on greenhouse gas emissions from high organic carbon grasslands. Biogeosciences 11: 6187–6207.

<http://www.biogeosciences.net/11/6187/2014/bg-11-6187-2014.html>

Eickenscheidt, T., Heinichen, J., Augustin, J., Freibauer, A. & M. Drösler, 2014: Nitrogen mineralization and gaseous nitrogen losses from waterlogged and drained organic soils in a black alder (*Alnus glutinosa* (L.) Gaertn.) forest. Biogeosciences 11: 2961–2976.

<http://www.biogeosciences.net/11/2961/2014/bg-11-2961-2014.html>

Eickenscheidt, T., Heinichen, J. & M. Drösler, 2015: The greenhouse gas balance of a drained fen peatland is mainly controlled by land-use rather than soil organic carbon content. Biogeosciences 12: 5161–5184.

<http://www.biogeosciences.net/12/5161/2015/bg-12-5161-2015.html>

Hommeltenberg, J., Mauder, M., Drösler, M., Heidbach, K., Werle, P. & H. P. Schmid, 2014. Ecosystem scale methane fluxes in a natural temperate bog-pine forest in southern Germany. Agricultural and Forest Meteorology 198–199: 273–284

<http://www.sciencedirect.com/science/article/pii/S0168192314002081>

Hommeltenberg, J., Schmid, H. P., Drösler, M. & P. Werle, 2014: Can a bog drained for forestry be a stronger carbon sink than a natural bog forest? Biogeosciences 11: 3477–3493.

<http://www.biogeosciences.net/11/3477/2014/bg-11-3477-2014.html>

Leiber-Sauheitl, K., Fuß, R., Voigt, C. & A. Freibauer, 2014: High CO<sub>2</sub> fluxes from grassland on histic Gleysol along soil carbon and drainage gradients. Biogeosciences 11: 749–761.

<http://www.biogeosciences.net/11/749/2014/bg-11-749-2014.html>

Leiber-Sauheitl, K., Fuß, R., Burkart, S., Buegger, F., Dänicke, S., Meyer, U., Petzke, K. J. & A. Freibauer, 2015: Sheep excreta cause no positive priming of peat-derived CO<sub>2</sub> and N<sub>2</sub>O emissions. *Soil Biology and Biochemistry* 88: 282-293.  
<http://www.sciencedirect.com/science/article/pii/S0038071715002060>

## Modellierung, Synthesen und Regionalisierung von Treibhausgasemissionen

Hoffmann, M., Jurisch, N., Albiac Borraz, E., Hagemann, U., Drösler, M., Sommer, M. & J. Augustin, 2015. Automated modeling of ecosystem CO<sub>2</sub> fluxes based on periodic closed chamber measurements: A standardized conceptual and practical approach. *Agricultural and Forest Meteorology* 200: 30–45.

<http://www.sciencedirect.com/science/article/pii/S0168192314002160>

Leppelt, T., Dechow, R., Gebbert, S., Freibauer, A., Lohila, A., Augustin, J., Drösler, M., Fiedler, S., Glatzel, S., Höper, H., Järveoja, J., Lærke, P. E., Maljanen, M., Mander, Ü., Mäkiranta, P., Minkkinen, K., Ojanen, P., Regina, K. & M. Strömgren, 2014: Nitrous oxide emission budgets and land-use-driven hotspots for organic soils in Europe. *Biogeosciences* 11: 6595–6612.  
<http://www.biogeosciences.net/11/6595/2014/bg-11-6595-2014.html>

Metzger, C., Jansson, P.-E., Lohila, A., Aurela, M., Eickenscheidt, T., Belotti-Marchesini, L., Dinsmore, K. J., Drewer, J., van Huissteden, J. & M. Drösler, 2015: CO<sub>2</sub> fluxes and ecosystem dynamics at five European treeless peatlands – merging data and process oriented modeling. *Biogeosciences* 12: 125–146.

<http://www.biogeosciences.net/12/125/2015/bg-12-125-2015.html>

Tiemeyer, B., Albiac Borraz, E., Augustin, J., Bechtold, M., Beetz, S., Beyer, C., Drösler, M., Eickenscheidt, T., Ebli, M., Fiedler, S., Förster, C., Freibauer, A., Giebels, M., Glatzel, S., Heinichen, J., Hoffmann, M., Höper, H., Jurasinski, G., Leiber-Sauheitl, K., Peichl-Brak, M., Roßkopf, N., Sommer, M. & J. Zeitz, 2016. High emissions of greenhouse gases from grasslands on peat and other organic soils. *Global Change Biology* 22: 4134–4149.  
<http://onlinelibrary.wiley.com/doi/10.1111/gcb.13303/abstract>

## Hydrologie und Bodenphysik

Bechtold, M., Tiemeyer, B., Laggner, A., Leppelt, T., Frahm, E. & S. Belting, 2014. Large-scale regionalization of water table depth in peatlands optimized for greenhouse gas emission upscaling. *Hydrology and Earth System Sciences* 18: 3319–3339.

<http://www.hydrol-earth-syst-sci.net/18/3319/2014/hess-18-3319-2014.html>

Dettman, U. & M. Bechtold, 2016. Deriving effective soil water retention characteristics from shallow water table fluctuations in peatlands. *Vadose Zone Journal* 15(10), doi:10.2136/vzj2016.04.0029  
<https://dl.sciencesocieties.org/publications/vzj/articles/15/10/vzj2016.04.0029>

Dettman, U. & M. Bechtold, 2016. One-dimensional expression to calculate specific yield for shallow groundwater systems with microrelief. *Hydrological Processes* 30(2): 334–340.  
<http://onlinelibrary.wiley.com/doi/10.1002/hyp.10637/abstract>

Dettmann, U., Bechtold, M., Frahm, E. & B. Tiemeyer, 2014. On the applicability of unimodal and bimodal van Genuchten-Mualem based models to peat and other organic soils under drying conditions. *Journal of Hydrology* 515: 103-115.  
<http://www.sciencedirect.com/science/article/pii/S0022169414003217>

## **Nährstoffe und gelöster organischer Kohlenstoff**

Frank, S., Tiemeyer, B., Gelbrecht, J. & A. Freibauer, 2014. High soil solution carbon and nitrogen concentrations in a drained Atlantic bog are reduced to natural levels by 10 years of rewetting. *Biogeosciences* 11: 2309–2324.  
<http://www.biogeosciences.net/11/2309/2014/bg-11-2309-2014.html>

Frank, S., Tiemeyer, B., Bechtold, M., Lücke, A. & R. Bol, 2017. Effect of past peat cultivation practices on present dynamics of dissolved organic carbon. *Science of the Total Environment* 574: 1243-1253.  
<http://www.sciencedirect.com/science/article/pii/S004896971631508X>

Tiemeyer, B. & P. Kahle, 2014. Nitrogen and dissolved organic carbon (DOC) losses from an artificially drained grassland on organic soils. *Biogeosciences* 11: 4123–4137.  
<http://www.biogeosciences.net/11/4123/2014/bg-11-4123-2014.html>

Tiemeyer, B., Pfaffner, N., Frank, S., Kaiser, K. & S. Fiedler, 2017. Pore water velocity and ionic strength effects on DOC release from peat-sand mixtures: Results from laboratory and field experiments. *Geoderma* 296: 86–97.  
<http://www.sciencedirect.com/science/article/pii/S0016706117302859>

## **Bodenkundliches und Karte organischer Böden**

Altdorff, D., Bechtold, M., van der Kruk, J., Vereecken, H. & J.A. Huisman, 2016: Mapping peat layer properties with multi-coil offset electromagnetic induction and laser scanning elevation data. *Geoderma* 261: 178-189.  
<http://www.sciencedirect.com/science/article/pii/S0016706115300252>

Fell, H., Rosskopf, N. & J. Zeitz, 2014: Impact of the spatial resolution of soils data on climate reporting for organic soils using the example of Germany. *Mires and Peat* 14: 1-9.  
<http://mires-and-peat.net/pages/volumes/map14/map1403.php>

Rosskopf, N., Fell, H. & J. Zeitz, 2015: Organic soils in Germany, their distribution and carbon stocks. *Catena* 133: 157-170.  
<http://www.sciencedirect.com/science/article/pii/S0341816215300126>